

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/107156/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Shirinfar, Bahareh and Ahmed, Nisar ORCID: <https://orcid.org/0000-0002-7954-5251> 2018. Chemical glycosylations for the synthesis of building units of post-translational modifications. *Helvetica Chimica Acta* 101 (1), e1700226. 10.1002/hlca.201700226 file

Publishers page: <http://dx.doi.org/10.1002/hlca.201700226>
<<http://dx.doi.org/10.1002/hlca.201700226>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Chemical Glycosylations for the Synthesis of Building Units of Post-translational modifications

Bahareh Shirinfar^{1,2} and Nisar Ahmed^{1,2,3*}

*E-mail: AhmedN14@cardiff.ac.uk

¹School of Chemistry, University of Bristol, Bristol, BS8 1TS, United Kingdom.

²Organic Chemistry Institute, University of Zürich, Winterthurerstrasse 190, 8057 Zürich, Switzerland

³School of Chemistry, Cardiff University, Cardiff, CF10 3AT, United Kingdom

Keywords: glycosylation, glycosidic bond, post-translational modification, amino acid, adenosine, riboside

Abstract

In chemical glycosylation reactions, a glycosyl donor couple with glycosyl acceptor through glycosidic linkage. Most of the products end up with mixture due to the formation of stereogenic center at anomeric carbon. Activation with suitable Lewis acid and introduction of the non-participating protecting group on donor and acceptor results in a selective product. Herein, we used suitably protected donor and acceptor which produced orthogonally protected building block with α -selectivity. We used also donor for the synthesis of modified phosphoribosylated amino acid. The formation of glycoside products can be used to synthesize complex biologically important organic molecules.

Introduction

The glycosylations are considered to play a vital role in a wide range of regulatory processes and glycoside products can be used to synthesize complex organic molecules which are of vital importance for the maintenance of diverse cellular functions.^[1] Poly-adenosine diphosphate (ADP) ribosylation is one of the complex polymeric post-translational modification (PTM) that occur at various functional groups in amino acid side chains (Fig. 1).^[2-5] The carbohydrates occur as polysaccharides, glycoconjugates or glycosides and in their chemical synthesis, the main challenge is to build glycosidic linkages between the monomeric units with appropriate orientation and most of glycosylations give a mixture of the product.^[6-9] The improvement of new methodologies for chemical glycosylation has emerged as an active area of research. Several types of new glycosyl donors for glycosylation reactions have been synthesized and applied. Diverse activation protocols have been established for the successful assembly of oligosaccharides from protected building blocks. Herein, we describe the glycosylation reactions resulted in synthesis of a ribosylated-amino acid **8**, and adenosine **9**. The glycosylation of adenosine gave an orthogonal bifunctional building block **9**, which may be used for future coupling reactions with amino acids and in the synthesis of PTMs (Fig. 1).






Figure 1. Post-translational protein modification (PTM): e.g; Poly-ADP-ribose polymer attached to protein.

We developed a synthetic route for the synthesis of benzyl protected phosphorylated ribofuranosyl *N*-phenyltrifluoroacetimidate **1** (Scheme 1) which was used in two representative glycosylation reactions; in synthesis of compounds **8** and **9**. The synthesis of **1** involved the following synthetic steps: i) synthesis of methyl D-ribofuranoside **3** according to the reported procedure,^[10] ii) phosphorylation of methyl D-ribofuranoside **3** through using P-amidite chemistry,^[11] iii) benzylation of phosphorylated methyl D-ribofuranoside **4** to protect the secondary hydroxyl groups,^[6,7] iv) deprotection of the anomeric methoxy to hydroxyl group via hydrolysis to give **6**, v) and finally introduction of the (*N*-phenyl)-2,2,2-trifluoroacetimido group on the anomeric hydroxy function group (Scheme 1).^[8,9]




Scheme 1. Synthesis of benzyl-phosphorylated ribofuranosyl *N*-phenyltrifluoroacetimidate **1**.

After the synthesis of donor **1**, 3,5-*O*-di-*tert*-butylsilyl protected *N*-benzoyl adenosine **2** was synthesized in order to obtain a bifunctional building block **9**. The synthesis of compound **2** was achieved according to a reported procedure.^[12-14] However, The phosphorylated donor **1** was also used for the preparation of ribosylated amino acid **8** (Scheme 3).^[15,16] After the C-terminal deprotection of **8** by using 20 mol% Pd(PPh₃)₄ and excess dimedone in THF, give building block **11** with a terminal phosphoribose moiety which can be used in future for coupling reactions and phosphorylations with precise control over the number of modifications.^[17,18]



Scheme 2. Synthesis of orthogonal bifunctional building blocks **9**, **10**.



Scheme 3. Synthesis of ribosyl amino acid derivatives **8**, **11**.

Generally, the synthesis of nucleosides with *O*-glycosylation on 2'-OH are considered difficult reactions, causing the formation of sideproducts in the reaction mixture and decreasing the yield of the desired *O*-glycosylated nucleoside.^[6-8] Some reports containing examples of the glycosylation of nucleosides with pyranosyl moieties, shimofuridin,^[19,20] and adenophostin,^[21-23] possessing 2'-*O*- β -fucopyranosyl and 3'-*O*- α -glucopyranosyl respectively have been reported. A number of synthetic nucleosides with *O*- β -linked furanosyl groups are also reported.^[24-26] However, very few examples have been reported for the synthesis of 2'-*O*- α -D-ribofuranosyladenosine which have 2'-*O*- α -ribosylated adenosine.^[15-18,27] In one example, the synthesis of 2'-*O*- α -D-ribofuranosyladenosine involved selective *trans*-coupling of arabinofuranose to adenosine, then subsequent attachment of the 2'-*O*-acyl group, proceeding by inversion of the stereochemistry at the 2-position of the arabinofuranose *via* an oxidation reduction sequence. To significantly improve previous procedures, we utilized the building block, di-*tert*-butylsilyl-protected adenosine nucleoside (**2**, Scheme 2).^[12-16] Using this block, and benzyl-phospho-ribofuranosyl *N*-phenyltrifluoroacetimidate as a donor **1**, it was possible to synthesize a α -riboside in excellent yield with orthogonal protective groups (Scheme 2). Filippov et al. reported similar kind of phosphorylated orthogonal building block in multiple steps synthesis (8-10 steps) using protecting and deprotecting chemistry and finished with laborious work up to provide free primary OH that was further functionalized with phosphoramidite group.^[16] We synthesized phosphorylated orthogonal building block with free primary OH in two steps which possibly be functionalized with phosphoramidite group.

The imidate **1** activated with TMSOTf (Lewis acid) in the presence of acceptor **2** resulted in the formation of **9** with excellent α -selectivity in 85% yield (Scheme 2). The stereochemistry was assigned according to the literature.^[28-30] The exocyclic amine of acceptor was dibenzoylated to avoid the side reaction. However, non-participating benzyl protection and (*N*-phenyl)-2,2,2-trifluoroacetimido group on the ribofuranosyl donor predominantly renders the α -product in glycosylation reactions. We deprotected silyl group of orthogonal bifunctional building block **9** to get compound **10** in 79% yield. However, the complete deprotection of **9** can be attempt by following the given literature.^[6,7] We also synthesized phosphoribosylated amino acid **8** in 73% yield (Scheme 3) by using imidate **1** and glutamic acid **7**. For this glycosylation reaction, also the TMSOTf used as an activator. The compounds **10** and **11** can possibly be used for extension of phosphate and peptide chain for PTMs.^[17,18]

Conclusions

We successfully synthesized ribosyl donor **1**, phosphoribosylated amino acid derivatives **8**, **11** and orthogonal bifunctional building blocks **9** and **10** which may provide an opportunity for the preparation of synthetic material for PTMs in a defined way.

Experimental Section

General

All chemicals were purchased from *Sigma-Aldrich* and *Merck*. Reactions were carried out under N₂ or Ar. Solvents for reactions were distilled prior to their use. Evaporation of the solvents *in vacuo* was done with the rotary evaporator. *Merck* silica gel 60 (40 – 63 μ m) was used for column chromatography. *Merck* thin layer chromatography plates silica gel 60 on aluminum were used for the spots visualizing by UV light. KMnO₄ and *p*-anisaldehyde solutions were used for staining. For ¹H-NMR spectra; *Bruker AV-300* (300 MHz), *Bruker AV-400* (400 MHz) and *Bruker AV-500* (500 MHz) were used. For ¹³C-NMR spectra; *Bruker AV-300* (75.5 MHz), *Bruker AV-400* (100 MHz) and *Bruker AV-500* (125 MHz) were used. For ³¹P-NMR spectra; *Bruker AV-300* (75.5 MHz) or *Bruker AV-400* (161 MHz) were used.

Experimental

Synthesis of 4. 4,5-Dicyanoimidazole (DCI, 1.95 g, 16 mmol) was added to a soln. of **3** (2.464 g, 15 mmol) and (BnO)₂PNiPr₂ (technical grade, 90%, 5.76 g, 15 mmol) in DMF (15 mL) at –10°C. The soln. was stirred at –10°C. After 5 h mCPBA (70%, 4.04 g, 18 mmol) was added in small portions and after stirring for another 0.5 h at 25°C the reaction soln was diluted with Et₂O and washed with H₂O (3 x 30 mL). The aq. layer was extracted with Et₂O (3 x 30 mL) and the combined org. layer were dried with MgSO₄. Evaporation of solvent *in vacuo* delivered colorless solid as crude product of **4** (7.37 g). Filtration over silica gel (CH₂Cl₂/MeOH 10:0 to 10:1) delivered colorless oil of **4** (3.61 g, 9 mmol, 60%). ¹H-NMR (400 MHz, CDCl₃): 7.41 – 7.30 (*m*, 2 x *Ph*), 5.05 – 5.03 (*m*, 2 x OCH₂Ph), 4.67 (*s*, 1 H), 4.17 – 4.09 (*m*, 1 H), 3.99 – 3.88 (*m*, 3 H), 3.74 (*d*, *J* = 4.1, C1H), 3.19 (*s*, CH₃), 2.89 (*s*, C3OH), 2.73 (broad *s*, C2OH). ¹³C-NMR (100 MHz, DMSO): 136.1, 136.0 (2*s*, 2 x OCH₂Ph), 128.5,

128.3, 127.8 (10*d*, 2 x OCH₂Ph), 108.3 (*d*, C1), 80.6 (*d*, C4), 74.0 (*d*, C2), 70.6 (*d*, C3), 3 x 68.5 (3*t*, C5, 2 x CH₂Ph), 54.3 (*q*, CH₃). ³¹P-NMR (161 MHz, CDCl₃): −3.44 (*quin*). HR-ESI-MS (MeOH + NaOH): 447.11782 (C₂₀H₂₅NaO₈P⁺, [M + Na]⁺, calc. 447.11793, Δ*m* = 0.2 ppm).

Synthesis of 5. NaH (60%, 0.64 g, 16 mmol) was added in small portions to a soln. of **4** (3.08 g, 7 mmol) in dry DMF (25 mL) at −10°C. Benzyl bromide (2.76 g, 16 mmol) was added after 5 min to the pale yellow and bubbling suspension. The reaction mixture was stirred for 1.5 h at −10°C before it was allowed to warm to 0°C. It was stirred for another 3.5 h at 0°C before TLC showed full consumption of **4** and the reaction was quenched by addition of H₂O (30 mL). The mixture was extracted with Et₂O (3 x 150 mL) and the collected org. layers were dried with MgSO₄. Evaporation of the solvent *in vacuo* delivered turbid pale yellow liquid as crude product of **5** (4.98 g). Purification by column chromatography (pentane/Et₂O 1:1 to 1:3) delivered colorless oil of **5** (3.36 g, 5 mmol, 56 %). ¹H-NMR (400 MHz, CDCl₃): 7.33 – 7.10 (*m*, 4 x Ph), 5.08 (*s*, 1 H), 2 x 4.93 (2*d*, *J* = 7.8, 2 x POCH₂Ph), 4.79 (*s*, 1 H), 4.52 (*q*, *J* = 12, 1 H), 4.42 (*d*, *J* = 11.7, 1 H), 4.32 (*d*, *J* = 11.7, 1 H), 4.27 – 4.16 (*m*, 1 H), 4.10 – 3.89 (*m*, C₅H₂, there in 3.99 – 3.96 (*t*-like *m*, C₄H)), 3.73 (*d*, *J* = 4.5, C₁H), 3.18 (*s*, CH₃). ¹³C-NMR (100 MHz, CDCl₃): 137.8 (4*s*, Ph), 128.7, 3 x 128.6, 128.3, 128.1, 2 x 128.0 (20*d*, 4 x Ph), 106.4 (*d*, C1), 79.8 (*d*, C4), 79.6 (*d*, C2), 78.1 (*d*, C3), 72.7, 72.5 (2*t*, 2 x COCH₂Ph), 2 x 69.4 (2*t*, 2 x POCH₂Ph), 68.1 (*t*, C5), 55.1 (*s*, CH₃). ³¹P-NMR (161 MHz, CDCl₃): −1.06 (*quin*). HR-ESI-MS (MeOH + NaOH): 627.21201 (C₃₄H₃₇NaO₈P⁺, [M + Na]⁺, calc. 627.21183, Δ*m* = 0.3 ppm).

Synthesis of 6. 5.5 ml HCl was added to a solution of **5** (1.80 g, 3 mmol) in CH₃CN (55 mL). The solution was stirred at room temperature for 2 h. Then, ethyl acetate (excess) and brine solution were added to quench reaction. Organic layer was separated and dried over Na₂SO₄ and concentrated *in vacuo*. The desired alcohol was purified by Silica gel column chromatography (CH₂Cl₂/EtOAc (50:1 to 50:10)) (1.20g, mmol, 60% yield) as a transparent oil. ¹H-NMR (400 MHz, DMSO): 7.37 – 7.05 (*m*, 20 H), 5.05 – 4.85 (*m*, 4 H), 4.66 – 4.34 (*m*, 4 H), 4.25 – 3.96 (*m*, 4 H), 3.91 – 3.68 (*m*, 3 H). ¹³C-NMR (100 MHz, CDCl₃) δ: 137.75, 135.62, 135.94, 135.90, 135.87, 135.83, 128.59, 128.46, 127.94 (arom), 106.31, 79.73, 77.98, 72.56, 69.34, 67.98. ³¹P-NMR (161 MHz, DMSO): −0.71 (*quin*). ESI-MS (MeOH + NaOH): 613.2 (C₃₅H₃₅NaO₈P⁺, [M + Na]⁺, calc. 613.2).

Synthesis of 1. Compound **6** (120 mg, 0.2 mmol) was dissolved in acetone (3 mL) and 100 μL water. Cs₂CO₃ (71 mg, 0.2 mmol, 1.1 eq.) and 2,2,2-trifluoro-*N*-phenylacetimidoyl chloride ClC(=NPh)CF₃ (30 μL, 0.2 eq.) were added and the reaction mixture was stirred at room temperature for 7 hours. After filtration over celite, the solvents were removed and the residue was purified using silica gel column chromatography (CH₂Cl₂/EtOAc (10:1) to afford the title compound as a colourless oil (107 mg, 70%). ¹H-NMR (400 MHz, CDCl₃) δ: 7.52 – 7.16 (*m*, 22H, arom), 7.02 (*m*, 1H, arom), 6.71 (dd, *J* = 16.7, 7.5 Hz, 2H, arom), 6.20 (*s*, 1H, H1), 4.95 – 4.91 (*m*, 4H, CH₂ Bn), 4.63 – 4.60 (*m*, 1H, H4') 4.54 – 4.33 (*m*, 4H, CH₂ Bn), 4.15 – 4.11 (*m*, 1H, H3'), 4.08 – 4.01 (*m*, 1H, H2'), 3.96 (AB, *J* = 11.3, 4.3 Hz, 1H, H5'), 3.85 (AB, *J* = 11.3, 4.0 Hz, 1H, H5'). ¹³C-NMR (100 MHz, CDCl₃) δ: 142.55,

136.17, 134.79, 134.67, 128.24, 127.72, 127.51, 127.11, 127.01, 126.91, 126.59, 125.15, 123.31, 119.57, 118.46, 101.11 (C1), 80.06 (C4), 77.25 (C2), 76.32 (C3), 71.77, 71.32, 68.38 (CH₂ Bn), 65.73 (C5). ³¹P-NMR (161 MHz, CDCl₃): -1.28. ESI-MS (MeOH + NaI): 762.8 (C₄₁H₃₉F₃NO₈P⁺, [M + H]⁺, calc. 761.7).

Synthesis of 9. The trifluoroacetimidate donor dissolved (**1**, 152mg, 0.2 mmol) in dry CHCl₃ (1ml), was added slowly at 0 °C to the solution of acceptor **2** (100mg, 0.16 mmol) and Trimethylsilyl trifluoromethanesulfonate (TMSOTf) (9ul, 0.2 eq.) in dry CHCl₃ (1 ml) containing freshly activated 4 Å molecular sieves, and then solution was stirred under Nitrogen atmosphere for 3h at 0 °C and quenched by the addition of triethylamine (TEA, 200 ul). The reaction mixture was concentrated *in vacuo*. After evaporation of the solvent the crude material was purified using silica gel column chromatography (hexane/EtOAc (3:2) as white foam (202 mg, 85% yield). ¹H NMR (500 MHz, CDCl₃) δ 8.58 (s, 1H, C2), 7.94 (s, 1H, C8), 7.88-7.25 (m, 30H, Arom. Bz, Bn), 6.06 (s, 1H, H1''), 5.40 (d, J = 4.0 Hz, 1H, H1'), 4.71 – 4.49 (m, 10H, CH₂ Bn's, C3'', C2''), 4.35-3.41 (m, 1H, C4'', C5'', C2', C3', C5'), 1.28 – 0.97 (m, 18H, t-butyl.). ¹³C NMR (125 MHz, CDCl₃) δ 170.10, 149.40, 142.87, 131.86, 131.06, 127.47, 126.78, 126.53, 126.37, 126.30, 126.22, 126.02, 125.89, 125.85, 125.78, 90.47 (C1'), 86.31 (C1''), 71.81, 71.05, 67.21, 60.89, 34.10, 31.40. ³¹P-NMR (161 MHz, CDCl₃): -1.17. ESI-MS (MeOH + NaI): 1189.5 (C₆₅H₇₁N₅O₁₃PSi⁺, [M + H]⁺, calc. 1188.4

Synthesis of 10. Compound **9** (120 mg, 0.1mmol) was dissolved in THF (5 mL). Treatment with TBAF (52.3mg, 2 eq.) at room temperature for one hour, concentration (Reaction was monitored by TLC, CH₂Cl₂: Methanol, 5:0.5) and purification using column chromatography (CH₂Cl₂: EtOAc, 3:4) afforded the title compound as a colourless syrup **10** (84 mg, 79% yield). ¹H NMR (500 MHz, CDCl₃) δ 8.51 (s, 1H, C2), 8.01 (s, 1H, C8), 7.94-7.14 (m, 30H, Arom. Bz, Bn), 6.36 (d, J = 7.4 Hz 1H, H1''), 5.91 (s, 1H, C5'' OH) 5.48 (d, J = 7.2 Hz, 1H, H1'), 5.19 4.71 – 4.49 (m, 4H, P-O-CH₂ Bn's) 4.91 – 4.80 (m, 5H, CH₂ Bn's, C4''), 4.75-4.32 (m, 5H, C2'', C3', C5'', C4') 4.28 (s, 1H, C3'' OH), 4.05-3.78 (m, 4H, C3'', C2', C5'). ¹³C NMR (125 MHz, CDCl₃) δ 173.10, 152.24, 145.58, 137.47, 136.38, 134.62, 133.91, 130.09, 129.42, 129.36, 129.26, 129.18, 128.99, 128.99, 128.85, 128.64, 128.45, 126.11, 102.09 (C1'), 100.11 (C1''), 71.32, 70.05, 66.29, 63.40, 56.10. ³¹P-NMR (161 MHz, CDCl₃): -1.24. ESI-MS (MeOH + NaI): 1048.35 (C₅₇H₅₅N₅O₁₃P⁺, [M + H]⁺, calc. 1048.35

Synthesis of 8. The trifluoroacetimidate donor dissolved (**1**, 38mg, 0.05 mmol) in dry CHCl₃ (0.5ml), was added slowly at 0 °C to the solution of glutamic acid **7** (20mg, 0.05 mmol) and Trimethylsilyl trifluoromethanesulfonate (TMSOTf) (2.5ul, 0.05 eq.) in dry CHCl₃ (0.5 ml) containing freshly activated 4 Å molecular sieves, and then solution was stirred under Nitrogen atmosphere for 6h at 0 °C and quenched by the addition of triethylamine (TEA, 200 ul). The reaction mixture was concentrated *in vacuo*. After evaporation of the solvent the crude material was purified using silica gel column chromatography (CH₂Cl₂/EtOAc (5:1) as white syrup (35 mg, 73% yield). ¹H-NMR (500 MHz, CDCl₃) δ: 7.67 – 7.17 (m, 29H, arom), 6.16 (s, 1H, H1'-α), 5.78 – 5.75 (m, 1H, -CH=), 5.78 – 5.75 (m, 1H, -CH=), 5.40-5.39 (d, 1H, =CH-), 5.23-3.22 (m, 19H, CH₂ Bn, -CH₂-, O-CH₂-, -, -CH-,

H3', H4', H5'), 2.40-1.92 , (m, 4H, -CH₂-CH₂-). ¹³C-NMR (125 MHz, CDCl₃) δ: 172.51, 171.28, 156.93, 144.01, 142.89, 138.14, 137.39, 136.15, 135.98, 132.07, 129.27-127.70 (arom.), 101.30, 100.11 (C1'), 99.35, 98.74, 95.57, 75.41, 73.63, 73.33, 73.08, 70.08, 66.69, 48.54, 47.77, 31.05, 27.22. ³¹P-NMR (161 MHz, CDCl₃): -0.97. ESI-MS (MeOH + NaI): 1004.35 (C₅₇H₅₅N₅O₁₃P⁺, [M + Na]⁺, calc. 1004.34)

Acknowledgments

Marie Skłodowska-Curie Actions COFUND (*Early Career Fellowship* 2016-2019, Marie Skłodowska-Curie grant agreement No 663830) and Novartis (Postdoctoral Fellowship 2013-2015) to Dr. Nisar. Ahmed are gratefully acknowledged. Royal Society Newton Postdoctoral Fellowship (2015-2017) to Dr. Bahareh shirinfar is acknowledged as well. The authors want to thank Henning J. Jessen (Albert-Ludwigs-University Freiburg, Germany) for his support at University of Zürich, Switzerland.

References:

1. T. Hunter, 'The Age of Crosstalk: Phosphorylation, Ubiquitination, and Beyond', *Mol. Cell.* **2007**, 28, 730 – 738.
2. P. O. Hassa, S. Hänni, M. Elser, M.O. Hottiger, 'Nuclear ADP-Ribosylation Reactions in Mammalian Cells: Where Are We Today and Where Are We Going?', *Microbiol. Mol. Biol. Rev.* **2006**, 70, 789 – 829.
3. S. Erener, V. Pétrilli, I. Kassner, R. Minotti, R. Castillo, R. Santoro, P. O. Hassa, J. Tschopp, M.O. Hottiger, 'Inflammasome-Activated Caspase 7 Cleaves PARP1 to Enhance the Expression of a Subset of NF-κB Target Genes', *Mol. Cell.* **2012**, 46, 200 – 211.
4. G. Asher, H. Reink, M. Altmeyer, M. Gutierrez-Arcelus, M.O. Hottiger, U. Schiebler, 'Poly(ADP-Ribose) Polymerase 1 Participates in the Phase Entrainment of Circadian Clocks to Feeding', *Cell* **2010**, 142, 943 – 953
5. D.A. Bobak, M. M. Bliziotis, M. Noda, S-C. Tsai, R. Adamik, J. Moss, 'Mechanism of Activation of Cholera Toxin by ADP-Ribosylation Factor (ARF): Both Low- and High-Affinity Interactions of ARF with Guanine Nucleotides Promote Toxin Activation', *Biochemistry* **1990**, 29, 855 – 861.
6. R. Das, B. Mukhopadhyay, 'Chemical O-Glycosylations: An Overview', *ChemistryOpen* **2016**, 5, 401 – 433.
7. G. J. Van der Heden van Noort, H. S. Overkleeft, G. A. van der Marel, D. V. Filippov, 'Ribosylation of Adenosine: An Orthogonally Protected Building Block for the Synthesis of ADP-Ribosyl Oligomers', *Org. Lett.* **2011**, 13, 2920 – 2923.

8. H. A. V. Kistemaker, G. J. Van der Heden van Noort, H. S. Overkleeft, G. A. van der Marel, D. V. Filippov, 'Stereoselective Ribosylation of Amino Acids', *Org. Lett.* **2013**, *15*, 2306 – 2309.
9. R. R. Schmidt, J. Michel, 'Facile Synthesis of α - and β -O-Glycosyl Imidates; Preparation of Glycosides and Disaccharides', *Angew. Chem. Int. Ed.* **1980**, *19*, 731–732.
10. N-S. Li, J. Lu, J. A. Piccirilli, 'Efficient Synthesis of Methyl 3,5-Di-O-benzyl- α -D-ribofuranoside and Application to the Synthesis of 2'-C- β -Alkoxymethyluridines', *Org. Lett.* **2007**, *9*, 3009-3012.
11. H. J. Jessen, N. Ahmed, A. Hofer, 'Phosphate esters and anhydrides – recent strategies targeting nature's favoured modifications', *Org. Biomol. Chem.* **2014**, *12*, 3526 – 3530.
12. X. F. Zhu, H. J. Williams Jr, A. I. Scott, 'An Improved Transient Method for the Synthesis of N-Benzoylated Nucleosides', *Synthetic Communications.* **2003**, *33*, 1233 – 1243.
13. T. Kamei, T. Fukaminato, N. Tamaoki, 'A photochromic ATP analogue driving a motor protein with reversible light-controlled motility: controlling velocity and binding manner of a kinesin–microtubule system in an in vitro motility assay', *Chem. Commun.* **2012**, *48*, 7625 – 7627.
14. R. A. Mizrahi, D. Shin, R. W. Sinkeldam, K. J. Phelps, A. Fin, D. J. Tantillo, Y. Tor, P. A. Beal, 'A Fluorescent Adenosine Analogue as a Substrate for an A-to-I RNA Editing Enzyme', *Angew. Chem. Int. Ed.* **2015**, *54*, 8713 –8716.
15. H. A. V. Kistemaker, A. P. Nardozza, H. S. Overkleeft, G. A. van der Marel, A. G. Ladurner, and D. V. Filippov, 'Synthesis and Macrodomein Binding of Mono-ADP-Ribosylated Peptides', *Angew. Chem. Int. Ed.* **2016**, *55*, 10634 –10638.
16. H. A. V. Kistemaker, L. N. Lameijer, N. J. Meeuwenoord, H. S. Overkleeft, G. A. van der Marel, D. V. Filippov, 'Synthesis of Well-Defined Adenosine Diphosphate Ribose Oligomers', *Angew. Chem. Int. Ed.* **2015**, *54*, 4915 – 4918.
17. G. S. Cremosnik, A. Hofer, H. J. Jessen, 'Iterative Synthesis of Nucleoside Oligophosphates with Phosphoramidites', *Angew. Chem. Int. Ed.* **2014**, *53*, 286 – 289.
18. S. Capolicchio, D. T. Thakor, A. Linden, H. J. Jessen, 'Synthesis of Unsymmetric Diphospho-Inositol Polyphosphates', *Angew. Chem. Int. Ed.* **2013**, *52*, 6912 – 6916.
19. J. Kobayashi, Y. Doi, M. Ishibashi, 'Shimofuridin A, a Nucleoside Derivative Embracing an Acylfucopyranoside Unit Isolated from the Okinawan Marine Tunicate *Aplidium multiplicatum*', *J. Org. Chem.* **1994**, *59*, 255 – 257.
20. J. Ning, Y. Xing, F. Kong, 'Efficient synthesis of a heptasaccharide, the repeating unit of the O-chain lipopolysaccharide produced by *Xanthomonas campestris* strain 642' *Carbohydr. Res.* **2003**, *338*, 1023 – 1031.
21. N. C. R. van Straten, G. A. van der Marel, J. H. van Boom, 'An Expeditious Route to the Synthesis of Adenophostin A', *Tetrahedron* **1997**, *53*, 6509 – 6522.

22. C. N. Borissow, S. J. Black, M. Paul, S. C. Tovey, S. G. Dedos, C. W. Taylor, B. V. L. Potter, 'Adenophostin A and analogues modified at the adenine moiety: synthesis, conformational analysis and biological activity', *Org. Biomol. Chem.* **2005**, 3, 245 – 252.
23. K. M. Sureshan, M. Trusselle, S. C. Tovey, C.W. Taylor, B. V. L. Potter, '2-Position Base-Modified Analogues of Adenophostin A as High-Affinity Agonists of the D-myo-Inositol Trisphosphate Receptor: In Vitro Evaluation and Molecular Modeling', *J. Org. Chem.* **2008**, 73, 1682 – 1692.
24. S. N. Mikhailov, E.V. Efimtseva, G.V. Gurskaya, M.V. Fomitcheva, S.V. Meshkov, 'An efficient synthesis and physico-chemical properties of 2'-O-D-ribofuranosylnucleosides, minor tRNA components', *J. Carbohydr. Chem.* **1997**, 16, 75 – 92.
25. E. V. Efimtseva, G. V. Bobkov, S. N. Mikhailov, A. Van Aerschot, G. Schepers, R. Busson, J. Rozenski, P. Herdewijn, 'Oligonucleotides Containing Disaccharide Nucleosides', *Helv. Chim. Acta* **2001**, 84, 2387 – 2397.
26. O. I. Andreeva, A. S. Golubeva, S. N. Kochetkov, A. Van Aerschot, P. Herdewijn, E. V. Efimtseva, B. S. Ermolinsky, S. N. Mikhailov, 'An Additional 20 -Ribofuranose Residue at a Specific Position of the DNA Primer Prevents Its Elongation by HIV-1 Reverse Transcriptase', *Bioorg. Med. Chem. Lett.* **2002**, 12, 681 – 684.
27. S. N. Mikhailov, I. V. Kulikova, K. Nauwelaerts, P. Herdewijn, 'Synthesis of 2'-O- α -D-ribofuranosyladenosine, monomeric unit of poly(ADP-ribose)' *Tetrahedron* **2008**, 64, 2871–2876.
28. H. Uchiro, T. Mukaiyama, 'A Significant Effect of a Lithium Salt in the Stereocontrolled Synthesis of α -D-Ribofuranosides' *Chem. Lett.* **1996**, 25, 271–272.
29. T. Mukaiyama, S. Suda, 'A New Class of S-Bridged Polynuclear Complex. The Formation and Crystal Structure of Spontaneously Resolved $[\{\text{Rh}(\text{aet})_3\}_4\text{Zn}_3\text{O}]^{4+}$ (aet = 2-aminoethanethiolate)', *Chem. Lett.* **1990**, 1143–1146.
30. C. H. Larsen, B. H. Ridgway, J. T. Shaw, D. M. Smith, K. A. Woerpel, 'Stereoselective C-Glycosylation Reactions of Ribose Derivatives: Electronic Effects of Five-Membered Ring Oxocarbenium Ions', *J. Am. Chem. Soc.* **2005**, 127, 10879 – 10884.

Table of Contents (TOC)

